

Table 5A-10. Number of black-crowned night herons detected during field surveys.

Habitat Type*	Iron Gate-Shasta	Iron Gate Reservoir	Fall Creek	Copco Bypass	Copco Reservoir	J.C. Boyle Peaking Reach	J.C. Boyle Bypass	J.C. Boyle Reservoir	Keno Canyon	Keno Reservoir	Link River	Total
<b><i>Plot Surveys</i></b>	<i>(n=18)</i>	<i>(n=38)</i>	<i>(n=16)</i>	<i>(n=4)</i>	<i>(n=37)</i>	<i>(n=72)</i>	<i>(n=22)</i>	<i>(n=20)</i>	<i>(n=18)</i>	<i>(n=23)</i>	<i>(n=18)</i>	<i>(n=286)</i>
Disturbed											1	1
Flyover										3	2	5
Lacustrine Unconsolidated Bottom										1		1
Riparian/Wetland Forest										5		5
Riparian/Wetland Scrub-shrub									8		3	11
Riverine Unconsolidated Bottom									1		1	2
<b><i>Facility Surveys</i></b>	<i>(n=1)</i>	<i>(n=3)</i>	<i>(n=4)</i>	<i>(n=3)</i>		<i>(n=1)</i>	<i>(n=2)</i>		<i>(n=1)</i>		<i>(n=3)</i>	<i>(n=18)</i>
All Habitats									5		6	11
<b>Total</b>									<b>14</b>	<b>9</b>	<b>13</b>	<b>36</b>

\*Detections were not recorded in habitat types not included in table.