

Table 5A-23. Number of yellow warblers detected during field surveys.

Habitat Type*	Iron Gate-Shasta	Iron Gate Reservoir	Fall Creek	Copco Bypass	Copco Reservoir	J.C. Boyle Peaking Reach	J.C. Boyle Bypass	J.C. Boyle Reservoir	Keno Canyon	Keno Reservoir	Link River	Total
<i>Plot Surveys</i>	(n=18)	(n=38)	(n=16)	(n=4)	(n=37)	(n=72)	(n=22)	(n=20)	(n=18)	(n=23)	(n=18)	(n=286)
Unidentified Habitat						1					2	3
Disturbed										2		2
LP							2					2
Montane Hardwood Oak		2			1	4	1					8
Montane Hardwood Oak-Conifer				1	1	6						8
Montane Hardwood Oak-Juniper		1	2			2						5
Mixed Chaparral						2			15			17
Pasture						3						3
Palustrine Emergent Wetland			1		1	1			2			5
Ponderosa Pine								2	2			4
Riparian/Wetland Forest	13	5		4	5	9				3	9	48
Riparian/Wetland Scrub-shrub	13	7	2		4	7	17	5	22	2	25	104
Riparian Grass	4					1			5	1	3	14
Riparian Mixed			1			6						7
Riverine Unconsolidated Bottom									1			1
<i>Facility Surveys</i>	(n=1)	(n=3)	(n=4)	(n=3)		(n=1)	(n=2)		(n=1)		(n=3)	(n=18)
All Habitats		1	1	1							6	9
<b>Total</b>	<b>30</b>	<b>16</b>	<b>7</b>	<b>6</b>	<b>12</b>	<b>42</b>	<b>20</b>	<b>7</b>	<b>47</b>	<b>8</b>	<b>45</b>	<b>240</b>

\*Detections were not recorded in habitat types not included in table.