

Table 5A-8. Avian TES species relative abundance per survey for all detections recorded during avian plot surveys.

(Birds detected per 20-minute survey period listed in order of decreasing relative abundance for all Project sections.)

<i>Bird Species</i>	Iron Gate-Shasta (n=18)	Iron Gate Reservoir (n=38)	Fall Creek (n=16)	Copco Bypass (n2=4)	Copco Reservoir (n=37)	J.C. Boyle Peaking Reach (n=72)	J.C. Boyle Bypass (n=22)	J.C. Boyle Reservoir (n=20)	Keno Canyon (n=18)	Keno Reservoir (n=23)	Link River (n=18)	Total (All Sections; n=286)
American White Pelican	0.00	1.39	0.00	0.00	0.54	0.11	0.04	1.40	0.44	6.09	4.06	1.16
Yellow Warbler	1.67	0.39	0.38	1.25	0.32	0.58	0.91	0.35	2.61	0.35	2.17	0.81
Forster's Tern	0.00	0.08	0.00	0.00	0.32	0.03	0.09	0.80	0.00	1.91	2.11	0.41
Acorn Woodpecker	0.06	0.34	0.31	0.25	1.65	0.38	0.00	0.05	0.00	0.00	0.00	0.38
Caspian Tern	0.00	0.58	0.00	0.00	0.46	0.00	0.05	0.35	0.06	0.39	0.11	0.21
Osprey	0.06	0.34	0.25	0.00	0.43	0.08	0.27	0.30	0.33	0.00	0.00	0.20
Great Egret	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.15	0.72	0.17	1.72	0.18
Vaux's Swift	0.06	0.00	0.06	0.00	0.19	0.43	0.18	0.10	0.00	0.00	0.00	0.16
Black-Crowned Night Heron	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.39	0.39	0.09
Yellow-Breasted Chat	0.67	0.00	0.06	0.00	0.05	0.10	0.00	0.00	0.00	0.00	0.00	0.08
Bald Eagle	0.00	0.05	0.00	0.00	0.22	0.06	0.05	0.20	0.00	0.09	0.00	0.07
Pileated Woodpecker	0.00	0.00	0.13	0.00	0.00	0.06	0.27	0.00	0.06	0.00	0.00	0.05
Willow Flycatcher	0.11	0.13	0.00	0.00	0.05	0.03	0.00	0.05	0.00	0.00	0.06	0.05
Pygmy Nuthatch	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.22	0.00	0.04
Lewis' Woodpecker	0.33	0.08	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.04
Olive-Sided Flycatcher	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.15	0.17	0.13	0.06	0.04
Black Tern	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.00	0.00	0.02
Black-Capped Chickadee	0.00	0.03	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.11	0.01
Mountain Quail	0.11	0.00	0.06	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.01
Northern Harrier	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.01
Western Bluebird	0.00	0.05	0.06	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Bufflehead	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.09	0.00	0.01
White-Faced Ibis	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.15	0.00	0.00	0.00	0.01
Blue-Gray Gnatcatcher	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Prairie Falcon	0.00	0.00	0.00	0.00	0.03	0.01	0.00	0.00	0.00	0.00	0.00	0.01
Common Loon	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 5A-8. Avian TES species relative abundance per survey for all detections recorded during avian plot surveys.

(Birds detected per 20-minute survey period listed in order of decreasing relative abundance for all Project sections.)

<i>Bird Species</i>	Iron Gate-Shasta (n=18)	Iron Gate Reservoir (n=38)	Fall Creek (n=16)	Copco Bypass (n2=4)	Copco Reservoir (n=37)	J.C. Boyle Peaking Reach (n=72)	J.C. Boyle Bypass (n=22)	J.C. Boyle Reservoir (n=20)	Keno Canyon (n=18)	Keno Reservoir (n=23)	Link River (n=18)	Total (All Sections; n=286)
Golden Eagle	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Snowy Egret	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.06	0.00	0.00	0.00
White-Headed Woodpecker	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00
Total (All Species)	3.06	3.58	1.31	1.75	4.32	1.90	1.95	4.80	4.94	10.00	10.78	4.08