

Table 5A-8. Avian TES species relative abundance per survey for all detections recorded during avian plot surveys.

(Birds detected per 20-minute survey period listed in order of decreasing relative abundance for all Project sections.)

<i>Bird Species</i>	Iron Gate-Shasta (n=18)	Iron Gate Reservoir (n=38)	Fall Creek (n=16)	Copco Bypass (n2=4)	Copco Reservoir (n=37)	J.C. Boyle Peaking Reach (n=72)	J.C. Boyle Bypass (n=22)	J.C. Boyle Reservoir (n=20)	Keno Canyon (n=18)	Keno Reservoir (n=23)	Link River (n=18)	Total (All Sections; n=286)
American White Pelican	0.00	1.39	0.00	0.00	0.54	0.11	0.04	1.40	0.44	6.09	4.06	<b>1.16</b>
Yellow Warbler	1.67	0.39	0.38	1.25	0.32	0.58	0.91	0.35	2.61	0.35	2.17	<b>0.81</b>
Forster's Tern	0.00	0.08	0.00	0.00	0.32	0.03	0.09	0.80	0.00	1.91	2.11	<b>0.41</b>
Acorn Woodpecker	0.06	0.34	0.31	0.25	1.65	0.38	0.00	0.05	0.00	0.00	0.00	<b>0.38</b>
Caspian Tern	0.00	0.58	0.00	0.00	0.46	0.00	0.05	0.35	0.06	0.39	0.11	<b>0.21</b>
Osprey	0.06	0.34	0.25	0.00	0.43	0.08	0.27	0.30	0.33	0.00	0.00	<b>0.20</b>
Great Egret	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.15	0.72	0.17	1.72	<b>0.18</b>
Vaux's Swift	0.06	0.00	0.06	0.00	0.19	0.43	0.18	0.10	0.00	0.00	0.00	<b>0.16</b>
Black-Crowned Night Heron	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.50	0.39	0.39	<b>0.09</b>
Yellow-Breasted Chat	0.67	0.00	0.06	0.00	0.05	0.10	0.00	0.00	0.00	0.00	0.00	<b>0.08</b>
Bald Eagle	0.00	0.05	0.00	0.00	0.22	0.06	0.05	0.20	0.00	0.09	0.00	<b>0.07</b>
Pileated Woodpecker	0.00	0.00	0.13	0.00	0.00	0.06	0.27	0.00	0.06	0.00	0.00	<b>0.05</b>
Willow Flycatcher	0.11	0.13	0.00	0.00	0.05	0.03	0.00	0.05	0.00	0.00	0.06	<b>0.05</b>
Pygmy Nuthatch	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.22	0.00	<b>0.04</b>
Lewis' Woodpecker	0.33	0.08	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	<b>0.04</b>
Olive-Sided Flycatcher	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.15	0.17	0.13	0.06	<b>0.04</b>
Black Tern	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00	0.00	0.00	<b>0.02</b>
Black-Capped Chickadee	0.00	0.03	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.11	<b>0.01</b>
Mountain Quail	0.11	0.00	0.06	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	<b>0.01</b>
Northern Harrier	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	<b>0.01</b>
Western Bluebird	0.00	0.05	0.06	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	<b>0.01</b>
Bufflehead	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.00	0.09	0.00	<b>0.01</b>
White-Faced Ibis	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.15	0.00	0.00	0.00	<b>0.01</b>
Blue-Gray Gnatcatcher	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	<b>0.01</b>
Prairie Falcon	0.00	0.00	0.00	0.00	0.03	0.01	0.00	0.00	0.00	0.00	0.00	<b>0.01</b>
Common Loon	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	<b>0.00</b>

Table 5A-8. Avian TES species relative abundance per survey for all detections recorded during avian plot surveys.

(Birds detected per 20-minute survey period listed in order of decreasing relative abundance for all Project sections.)

<i>Bird Species</i>	<b>Iron Gate-Shasta (n=18)</b>	<b>Iron Gate Reservoir (n=38)</b>	<b>Fall Creek (n=16)</b>	<b>Copco Bypass (n2=4)</b>	<b>Copco Reservoir (n=37)</b>	<b>J.C. Boyle Peaking Reach (n=72)</b>	<b>J.C. Boyle Bypass (n=22)</b>	<b>J.C. Boyle Reservoir (n=20)</b>	<b>Keno Canyon (n=18)</b>	<b>Keno Reservoir (n=23)</b>	<b>Link River (n=18)</b>	<b>Total (All Sections; n=286)</b>
Golden Eagle	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	<b>0.00</b>
Snowy Egret	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.06	0.00	0.00	<b>0.00</b>
White-Headed Woodpecker	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	<b>0.00</b>
<b>Total (All Species)</b>	<b>3.06</b>	<b>3.58</b>	<b>1.31</b>	<b>1.75</b>	<b>4.32</b>	<b>1.90</b>	<b>1.95</b>	<b>4.80</b>	<b>4.94</b>	<b>10.00</b>	<b>10.78</b>	<b>4.08</b>