

Table 7A-13. Number of aquatic fur-bearing mammals detected during relicensing field studies.

Species	Habitat Type*	<i>Iron Gate-Shasta</i>	<i>Iron Gate Reservoir</i>	<i>Fall Creek</i>	<i>Copco Bypass</i>	<i>Copco Reservoir</i>	<i>J.C. Boyle Peaking Rch.</i>	<i>J.C. Boyle Bypass</i>	<i>J.C. Boyle Reservoir</i>	<i>Keno Canyon</i>	<i>Keno Reservoir</i>	<i>Link River</i>	Total
Beaver	Riparian/Wetland Forested		13									1	14
	Riparian/Wetland Scrub-shrub	2		1									3
	Palustrine Unconsolidated Bottom			1			1		1		1		4
Mink	Riparian Grassland											4	4
	Riparian/Wetland Scrub-shrub									3			3
	Riverine Unconsolidated Bottom									2	1		3
	Palustrine Emergent Wetland					1							1
Muskrat	Lacustrine Unconsolidated Bottom			1						1	2		4
	Riparian Grassland										1		1
River Otter	Riverine Unconsolidated Shoreline	1					1	4					6
	Palustrine Unconsolidated Bottom						1						1
	Total	3	13	3	0	1	3	4	1	6	5	5	44

*Detections were not recorded in habitat types not included in table.