

Table 5A-16. Number of Forster's terns detected during field surveys.

Habitat Type*	Iron Gate-Shasta	Iron Gate Reservoir	Fall Creek	Copco Bypass	Copco Reservoir	J.C. Boyle Peaking Reach	J.C. Boyle Bypass	J.C. Boyle Reservoir	Keno Canyon	Keno Reservoir	Link River	Total
Plot Surveys	(n=18)	(n=38)	(n=16)	(n=4)	(n=37)	(n=72)	(n=22)	(n=20)	(n=18)	(n=23)	(n=18)	(n=286)
Flyover		1			3		3			3	13	23
Lacustrine Unconsolidated Bottom		2			9			14		36	8	69
Pasture										4		4
Ponderosa Pine								2				2
Riparian Grass										1		1
Riverine Unconsolidated Bottom						2					17	19
Facility Surveys	(n=1)	(n=3)	(n=4)	(n=3)		(n=1)	(n=2)		(n=1)		(n=3)	(n=18)
All Habitats							2		3		17	22
Reservoir Surveys		(n=4)			(n=4)			(n=2)		(n=4)	(n=1)	(n=15)
All Habitats		10						66		56		132
Total		13			12	2	5	82	3	100	55	272

*Detections were not recorded in habitat types not included in table.