

Table 7A-9. Relative abundance of six common bird species captured per 1,000 net hours by 10-day period during 1997-2000 at the Frain Ranch (TOPS) constant effort mist netting station. Ten-day periods start on May 1 and end October 15 (e.g., Period 3 = May 21 through 30 and Period 13 = August 28 through September 6).

Species	10-Day Time Periods															
	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Common Bushtit	0.00	0.00	0.00	0.00	13.33	0.00	0.00	0.00	0.00	0.00	35.12	0.00	7.10	33.33	25.15	0.00
Hammond's Flycatcher	8.11	0.00	0.00	0.00	0.00	0.00	0.00	6.63	0.00	6.56	10.04	4.43	21.31	0.00	0.00	0.00
Hermit Thrush	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.43	99.43	79.17	75.45	60.00
Lincoln's Sparrow	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	10.04	8.86	35.51	25.00	30.18	0.00
Orange-Crowned Warbler	8.11	9.83	0.00	0.00	0.00	0.00	21.43	6.63	20.00	13.11	15.05	17.72	28.41	4.17	10.06	0.00
Swainson's Thrush	8.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	25.09	4.43	0.00	4.17	10.06	0.00